arXiv:1403.4351v1 [gr-qc] 18 Mar 2014

The small and large D limit of Parikh-Wilczek tunneling model for Hawking radiation

June-Yu Wei[1,*] and Wen-Yu Wen[1,2,†]

[1]*Department of Physics and Center for High Energy Physics, Chung Yuan Christian University, Chung Li City, Taiwan*

[2]*Leung Center for Cosmology and Particle Astrophysics National Taiwan University, Taipei 106, Taiwan*

Abstract

In this letter, we study both the small and large dimension D limit of the tunneling model of Hwaking radiation proposed by Parikh and Wilczek[1]. We confirm that the result $\Gamma \sim e^{\Delta S}$ is still valid for arbitrary $D > 3$ and universal prefactor 8π in the area spectrum. The sensible large D limit is given by $\sqrt{D} \ll r_0$ in order to have nonzero radiation. On the other hand, the sensible small D limit is given by taking $D = 3+\epsilon$ as a continuous parameter. We also explicitly show the leading order correction to the thermal radiation.

*E-mail: weijuneyu@gmail.com

†E-mail: steve.wen@gmail.com

I. INTRODUCTION

It was argued and shown in [2] that in the limit $N \to \infty$, General Relativity dramatically simplifies and $1/N$ could serve as a useful expansion parameter. Dynamics of a black hole with the horizon r_0 is controlled by two relevant small scales: $r_0/\sqrt{D}$ due to horizon area and r_0/D form the strong localization of gravitational potential. In this paper, we would like to confirm prediction from the tunneling model of Hawking radiation proposed by Parikh and Wilczek and study its nonthermal correction in the limits of large and small D. We start with a brief review of the D-dimensional black hole metric and its thermal quantities. A D-dimensional Schwarzschild black hole metric is given in the Schwarzschild-Tangherlini form[3]:

$$\begin{aligned} ds^2 = \quad & -f(r)dt_s^2 + f(r)^{-1}dr^2 + r^2 d\Omega_{D-2}^2, \\ & f(r) = 1 - (\frac{r_0}{r})^{D-3} \end{aligned} \tag{1}$$

where $r = r_0$ is the horizon such that $r_0^{D-3} = \frac{16\pi}{(D-2)\Omega_{D-2}}M$. The volume of unit sphere $\Omega_{D-2} = \frac{2\pi^{(D-1)/2}}{\Gamma(\frac{D-1}{2})}$.The Hawking temperature is given by

$$T_H = \frac{D-3}{4\pi r_0}. \tag{2}$$

The apparent singularity at the horizon can be removed by a coordinate transformation of the Painlevè-type:

$$t = t_s + r_0^{(D-3)/2} \int \frac{dr}{\sqrt{r^{D-3} - r_0^{D-3}}}, \tag{3}$$

which brings into the metric suitable for our purpose:

$$ds^2 = -(1 - (r_0/r)^{D-3})dt^2 + 2(\frac{r_0}{r})^{(D-3)/2} dtdr + dr^2 + r^2 d\Omega_{D-2}^2. \tag{4}$$

We remark that the tunneling process across the horizon happens in this stationary vacuum, instead of static one given by metric (1).

II. TUNNELING MODEL IN D DIMENSIONS

In this section, we apply the Parikh-Wolczek tunneling model[1] to D dimensional black hole and obtain the general expression for tunneling rate in D dimensions. We regard

radiation as a shell of energy ω moving in the geodesics of a spacetime with M replaced by $M-\omega$ due to back reaction, where the radial null geodesics are given by

$$\dot{r} \equiv \frac{dr}{dt} = \pm 1 - \sqrt{(\frac{r_\omega}{r})^{D-3}}. \tag{5}$$

r_ω denotes the horizon after radiation, such that $r_\omega^{D-3} = \frac{16\pi}{(D-2)\Omega_{D-2}}(M-\omega)$. The WKB approximation states that the emission rate $\Gamma \sim e^{-2ImS}$, where the imaginary part of action reads

$$ImS = Im\int_{r_{in}}^{r_{out}} p_r dr = Im\int_{M}^{M-\omega}\int_{r_0}^{r_\omega}\frac{dH}{\dot{r}}dr, \tag{6}$$

for Haniltonian $H = M-\omega'$. The half line contour integral in the complexified r-plane picks up a residue at $r = r_{\omega'}$, that is

$$\int_{r_0}^{r_\omega}\frac{dr}{\dot{r}} = \frac{2\pi i}{D-3}\big(\frac{16\pi}{(D-2)\Omega_{D-2}}(M-\omega')\big)^{1/(D-3)}. \tag{7}$$

After substituting this result into integral (6), one obtains

$$-2ImS = \frac{4\pi}{D-2}\big(\frac{16\pi}{(D-2)\Omega_{D-2}}\big)^{\frac{1}{D-3}} M^{\frac{D-2}{D-3}}\big((1-\frac{\omega}{M})^{\frac{D-2}{D-3}} - 1\big) = \Delta S_{BH}, \tag{8}$$

which is exact the change of Bekenstein-Hawking entropy after radiation of energy ω for a D-dimensional Schwarzschild black hole. This explicit calculation illustrates that the Parikh-Wilczek tunneling model of Hawking radiation in arbitrary dimension $D > 3$ predicts the same result, that is $\Gamma \sim e^{\Delta S_{BH}}$.

III. AREA SPECTRUM IN D DIMENSIONS

If we consider the energy of black hole is quantized like any confined quantum system, then area or entropy quantization is also expected[5]. The quantization condition can be easily realized in (8) under the assumption that black hole energy is reduced by ΔM after each emission of quanta h, that is

$$-2ImS = h = 2\pi, \tag{9}$$

where we always use the natural units. Under the large black hole limit $\Delta M \ll M$, one obtains the quantization of black hole mass $\Delta M = hT_H$. On the other hand, one can show the differential relation between r_0 and M gives rise to $\Delta r = \frac{1}{2M}$. Therefore the horizon area is quantized as

$$\Delta A = 16\pi M \Delta r = 8\pi. \tag{10}$$

Our result confirms the area quanta takes a universal prefactor 8π as first shown in the [6] and many other kinds of black holes.

IV. LARGE D CORRECTION TO THE EMISSION RATE

In this section, we take large D limit of emission rate to see its correction to thermal spectrum. For $\omega \ll M$, we can expand

$$\Delta S \simeq -\frac{4\pi}{D-3} r_0 \omega + \frac{4\pi}{(D-3)^2} r_0 \frac{\omega^2}{M} + \cdots \tag{11}$$

The first term is nothing but Hawking's thermal radiation. We see that naively sending $D \to \infty$ corresponds to a black hole at infinite temperature. If at the same time keeping r_0/D finite, it corresponds to that at finite temperature. Both limits give rise to a trivial emission rate thanks to vanishing first term in (11) for sufficiently small ω. On the other hand, the sensible limit which keeps the first term finite while $\omega \ll M$ is given by that $r_0/\sqrt{D} \gg 1$. It was pointed out in [2] that the length scales like $r_0/\sqrt{D}$ in D dimensions, so this nontrivial limit asks for a black hole of sizable area to ensure a significant radiation. The emission rate after receiving nonthermal corretion from the second term in (11) reads $\Gamma = e^{(\sqrt{D}/r_0)^D}\Gamma_0$, where the usual Hawking thermal radiation $\Gamma_0 \equiv e^{-\omega/T_H}$.

V. SMALL D CORRECTION TO THE EMISSION RATE

It is also interesting to investigate (11) for small dimensions, say $D \to 3^+$[4]. To do that, we let $D = 3+\epsilon$ and then send $\epsilon \to 0$. This limit gives rise to zero temperature and no radiation. For nonvanishing radiation, one also needs $r_0^{1+\epsilon} \gg \epsilon$, while the emission rate after receiving nonthermal corretion from the second term in (11) reads $\Gamma \simeq e^{r_0^{-1-\epsilon}}\Gamma_0$.

VI. DISCUSSION

We have demonstrated that the tunneling model of Hawking radiation predicts the same result $\Gamma \sim e^{\Delta S}$ in arbitrary dimension D and confirm the universal prefactor 8π in area quantization. In particular, one can fix the scale r_0/D for nonvanishing Hawking temperature, or $r_0/\sqrt{D}$ for nonzero Hawking radiation. The nonthermal correction is also obtained in terms of D. A different way of taking large D limit of black hole was studied in the dimensionally continued gravity[7], where the blacken function in the metric (1) is replaced by $f(r) = 1 + \frac{r^2}{L^2} - (\frac{r_0}{r})^{\frac{2(D-D_c)}{(D_c-1)}}$, for some critical dimension $D_c \geq 3$. In particular, in the limit $D, L \to \infty$ but keepingD/D_c fixed, the solution becomes asymptotically flat

and temperature is finite. Due to the higher-curvature terms in the action, the Bekenstein-Hawking area law is disobeyed and $S \propto A^{\frac{D-D_c}{D}}$. Nevertheless, we still find that the result $\Gamma \sim e^{\Delta S}$ is valid. Our observation is that the Parikh-Wilczek tunneling model predicts the same result as long as the first law of thernodynamics is satisfied[1]. This reflects the fact that the tunneling model respects conservation of energy, which is the very spirit of the first law.

Acknowledgments

We are grateful to the help of Ji-Hsien Dai at the early stage of this work. This work is supported in parts by the Taiwan's National Science Council (grant No. 102-2112-M-033-003-MY4) and the National Center for Theoretical Science.

---

[1] M. K. Parikh and F. Wilczek, "Hawking radiation as tunneling," Phys. Rev. Lett. **85**, 5042 (2000)

[2] R. Emparan, R. Suzuki and K. Tanabe, "The large D limit of General Relativity," JHEP **1306**, 009 (2013) [arXiv:1302.6382 [hep-th]].

[3] F. R. Tangherlini, "Schwarzschild field in n dimensions and the dimensionality of space problem," Nuovo Cim. **27**, 636 (1963)

[4] V. Asnin, D. Gorbonos, S. Hadar, B. Kol, M. Levi and U. Miyamoto, "High and Low Dimensions in The Black Hole Negative Mode," Class. Quant. Grav. **24**, 5527 (2007) [arXiv:0706.1555 [hep-th]].

[5] S. Hod, "Bohr's correspondence principle and the area spectrum of quantum black holes," Phys. Rev. Lett. **81**, 4293 (1998) [gr-qc/9812002].

[6] M. Maggiore, "The Physical interpretation of the spectrum of black hole quasinormal modes," Phys. Rev. Lett. **100**, 141301 (2008) [arXiv:0711.3145 [gr-qc]].

[7] G. Giribet, "Large D limit of dimensionally continued gravity," Phys. Rev. D **87**, no. 10, 107504 (2013) [arXiv:1303.1982 [gr-qc]].

---

[1] In this particular limit, the black hole mass $M \sim \frac{r_0^{D-D_c}}{2G}$, Hawking temperature $T \sim \frac{\hbar(D-D_c)}{2\pi D_c r_0}$ and Bekenstein-Hawking entropy $S \sim \frac{\pi D_c r_0^{D-D_c}}{\hbar G(D-D_c)}$[7]. It is straightforward to check that the first law of thermodynamics $M = TS$ holds.